

Beach Morphology using Ground Penetrating Radar

Harry M. Jol, Department of Geography, University of Wisconsin - Eau Claire, 105 Garfield Ave., Eau Claire, WI; 54702, USA., jolhm@uwedc.edu

Curt Peterson, Department of Geology, Portland State University, PO Box 751, Portland, OR, 97297-0751, USA, curt@ch1.ch.pdx.edu

Peter Ruggerio, Washington Department of Ecology, P.O. Box 47690, Olympia, WA, 98504-7690, USA, prug461@ecy.wa.gov

Sandy Vanderburgh, Department of Geography, University-College of the Fraser Valley, Abbotsford, BC, V2S 4N2 CANADA, vanderburghs@ucfv.bc.ca

Jim Phipps, Department of Geology, Grays Harbor College, Aberdeen, WA 98520-7599, USA, jhipps@ghc.ctc.edu

Several ground penetrating radar (GPR) datasets that were requested by the participants of the research group were collected during the summer of 1998. 100 MHz topographically corrected GPR lines were surveyed to fill in gaps of previous data collection (1996 and 1997), particularly in the north section of the Columbia River Littoral Cell. The northern lines confirmed our hypothesis that deposition in the northern portion of the cell has not been occurring for long period (as compared to lower portions of the cell).




Figure 1 *Upper:* 100 MHz GPR line shot in Moclips – northern portion of the cell. Paleo-scarp located at 100-110 m. *Lower:* 100 MHz GPR line shot as a continuation of a 1997 line to confirm paleo-beach cliff location and lower Holocene – Pleistocene contact (35-55 m)

Field experiments that compared radar frequencies and transmitter powers were carried out at several locations within the cell. Antennae frequencies tested were 25, 50, 100 and 200 MHz. The transmitter powers used were 400 and 1000 volt (maximum allowable by federal government regulations). All lines were topographically corrected. The experiment allowed for both higher resolution datasets to be collected so that better insight into the sedimentological structures could be achieved, as well as deeper penetration, particularly into the large dunes of the Clatsop area of the study.




Figure 2 A frequency comparison of 3 GPR antennae frequencies. The site was located on the south end of Ocean Shores. The profiles are plotted with similar parameters. *Upper:* 200 MHz antennae – note higher resolution stratigraphy, *Middle:* 100 MHz antennae – note deeper penetration and lower resolution, *Lower:* 50 MHz antennae – note much deeper penetration (off this scale) but a loss of much of the near surface stratigraphy.

Two three-dimensional grids were collected, one in the North Grays Harbor area and one in the Long Beach area. The datasets were collected to image recent paleo-beachface (shoreface) deposits and compare the results to present beachfaces. A GPS grid was collected at the same time as the high resolution (200 MHz) GPR grid was collected. The dimension of the grid was 25 m by 25 m with datapoints collected every 0.5 m (0.5 m antennae step along lines separated by 0.5 m). Initial processing and interpretation reveals several paleo-beach layers that will be compared to the present GPS beachface surveys.




Figure 2 200 MHz 3D Grid shot to map paleo-beachfaces. Grid shown is located in the Long Beach portion of the cell. *Upper Left:* Three dimensional perspective of the dataset (red lines indicating shorefaces). *Upper Right:* One of the 51 GPR lines shot – each dipping reflection is interpreted as a paleo-beachface, *Lower:* The cube shown here has been cut away to show the internal stratigraphy of the shoreface. Note the continuity of the beachfaces.

Acknowledgements

The major support for this project comes from the Southwest Washington Coastal Erosion Study - United States Geological Survey (USGS). Further support comes from Sensors and Software, University of Wisconsin-Eau Claire, Portland State University, University College of the Fraser Valley and Grays Harbor College. We would also like to acknowledge the support of the Washington Department of Ecology (DOE), and Grays Harbor, Pacific, and Clatsop counties. Able bodied field assistance was provided by Brian Thayer, Mark Newman-Bennet, David Qualman, April Herb, Chad Bartz, Lorraine Woxel, Nick Zerr and Andrew Zachery (your long days in the field are appreciated!). We thank Rilea Armed Forces Training Center, the Cranberry Research Foundation, and the communities within the study area for their aid in this project.

References

- Atwater, B.F., 1996. Coastal evidence for great earthquakes in Western Washington. In *Assessing Earth Hazards and Reducing Risk in the Pacific Northwest*, US Geological Survey Professional Paper 1560, pp. 77-90, 1996.
- Jol, H.M., Peterson, C.D., Vanderburgh, S. and Phipps, J. 1998. GPR as a regional geomorphic tool: shoreline accretion/erosion along the Columbia river littoral cell. *Proceedings of the Seventh International Conference on Ground Penetrating Radar (GPR'98)*, May 27-30, University of Kansas, Lawrence, KS, USA, Volume 1, pp. 257-262.
- Jol, H.M., D.G. Smith, R.A. Meyers, and D.C. Lawton, 1996a. Ground penetrating radar: high resolution stratigraphic analysis of coastal and fluvial environments. In: *Stratigraphic Analysis using Advanced Geophysical, Wireline and Borehole Technology for Petroleum Exploration and Production*, J.A. Pacht, R.E. Sheriff, and B.F. Perkins, (eds.), Gulf Coast Section Society of Economic Paleontologists and Mineralogists Foundation 17th Annual Research Conference, pp. 153-163.
- Jol, H.M., D.G. Smith, and R.A. Meyers, 1996b. Digital ground penetrating radar (GPR): An improved and very effective geophysical tool for studying modern coastal barriers (examples for the Atlantic, Gulf and Pacific coasts, U.S.A.). *Journal of Coastal Research*, 12(4): 960-968.
- Kaminsky, G.M., P. Ruggiero, G. Gelfenbaum, and C. Peterson, 1997. Long term coastal evolution and regional dynamics of a US Pacific Northwest Littoral Cell, *Coastal Dynamics'97*.
- Meyers, R., D.G. Smith, H.M. Jol, and C.R. Peterson, 1996. Evidence for eight great earthquake-subsidence events detected with ground-penetrating radar, Willapa barrier, Washington. *Geological Society of America Geology*, 24: 99-102.
- Phipps, J.B., 1990. Coastal accretion and erosion in Washington State 1977-1987 Shorelands and Coastal Zone Management Program. Washington Department of Ecology, Olympia, Washington.